

Isolation, Purification and Characterisation of Proteins from Plants

In vivo study of proteins

Cellular localization of protein



Activity e.g. detection of metal transport by specific fluorescent dyes or patch clamping

Limitations :

No structural determination

No analysis of binding constant

No analysis of catalytic mechanism



Kobae et al., 2004, PlantCell Physiol 45, 1749-58



Leitenmaier B, Küpper H (2011) Plant Cell & Environment 34, 208-219

I) Protein Isolation: Sources (expression) of plant proteins

Native expression (i.e. in intact organs or in tissue culture)

➤ Advantages:

- Correct folding
- Correct post-translational processing (e.g. insertion of active centre, phosphorylation, glycosylation, cleavage, ...)
- No cloning needed

➤ Disadvantages

- Usually lower yield
- Often many similar proteins in the same organisms → difficulties with purification



I) Protein Isolation: Sources (expression) of plant proteins

Heterologous expression (e.g. in bacteria, yeast or insect cells)

➤ Advantages:

- Usually higher yield
- Specific expression of one protein in large amounts and possibly with specific tag (e.g. His-tag) facilitates purification

➤ Disadvantages

- Often problems with folding, in particular in the case of large proteins and integral membrane proteins
- Insertion of complicated cofactors (e.g. iron-sulfur clusters) is difficult or impossible, other post-translational modifications may be missing or different compared to the native protein → possibly wrong conclusions about functions/mechanisms

I) Protein Isolation: Methods for isolation

Grinding in liquid nitrogen

- Used for hard tissues and cells like roots, stems, but also for hard-walled cells like some algae and cyanobacteria
- Low temperature protects the proteins during grinding
- Time consuming (manual grinding) or requiring suitable machinery (expensive solution: lab mill, several thousand €, cheap solution: wheat mill, about 200 €)




Liquid nitrogen cooled
wheat mill in Küpper lab

I) Protein Isolation: Methods for isolation

Ultrasound homogenization ... Can cause artefacts of denaturation

Many peaks of metalloproteins were strongly reduced and/or moved to low MW, showing severe protein denaturation, already by „gentle“ ultrasound treatment of protein extracts


Küpper H, Hussain Bokhari SN, Jaime Perez N, Lyubenova L, Ashraf N, Andresen E (2019) Analytical Chemistry 91, 1710961-10969



I) Protein Isolation: Methods for isolation

French Press

- Used for individual cells (plant cell culture, algae or bacteria) without or with soft walls
- Does not require freezing and thus may avoid artefacts of freezing
- Requires very expensive (usually many thousand €) machinery



From: www.diversified-equipment.com




From: www.pegasusscientific.com

I) Protein Isolation: Methods for isolation

Lysis buffer

- Used only for bacteria or animal cells
- May cause degradation
- No machinery needed



From: bio-ggs.blogspot.com/2009/11/ggs-live-western...

II) Protein Purification: Overview of Principles

Separation by expression site

- Selective use of tissues or organelles
- Separation of soluble from membrane proteins by centrifugation

Separation by size

- Ultrafiltration
- Size exclusion chromatography
- Preparative native gel electrophoresis

Separation by charge

- Ion exchange chromatography
- Isoelectric focussing (as chromatography, in solution or in gel electrophoresis)

Separation by specific binding sites

- Metal affinity chromatography using natural or artificial metal binding sites
- Immuno-Chromatography using immobilised antibodies
- Magnetic separation using magnetically tagged antibodies

II) Protein Purification: Separation by size in native gels

- Principle: Small proteins are less retained by the fibers of the gel than large proteins, so that small proteins migrate **faster**





foto of a green gel (Küpper et al., 2003, Funct Plant Biol)

From: www.columbia.edu/.../c2005/lectures/lec6_09.html

III) Protein Identification: Overview of Principles

Size determination

- Size exclusion chromatography or SDS PAGE
- Comparison with expected size of protein (known e.g. from reference or cDNA)

Western Blotting

- Binding to specific primary antibody, detected via labelled or enzymatically active secondary antibody

Biochemical assays in native gels

- Identification of enzymes by their characteristic activity
- Identification of metalloproteins by their metal content

Mass Spectrometry


- Fragmentation of the protein, identification of fragment sizes, and subsequent comparison to a library of known fragmentation patterns

N-terminal Sequencing (Edman degradation)

- Sequential chemical removal of individual amino acids from the N-terminus

III) Protein Identification by assays in native gels


- Example of a biochemical assay in native gels: identification of Cd-Carboanhydrase by its metal content combined with an in-gel carboanhydrase assay



from: Lane TW, Morel FMM (2000) PNAS97, 4627-4631

III) Protein Identification by HPLC-ICPMS

- Example of a metalloprotein identification by HPLC-ICPMS → Cd binding to main light harvesting antenna LHCII




Andresen E, Kappel S, Stärk HJ, Riegger U, Borovec J, Mattusch J, Heinz A, Schmelzer CEH, Matoušková Š, Dickinson B, Küpper H (2016) New Phytologist 210, 1244-1258.

III) Protein Identification: Mass Spectrometry

Principle

- 1) protein band is cut out of gel
- 2) protein is digested into peptides
- 3) mass spectrometry
- 4) comparison of the fragment sizes with a database
- 5) assignment of likely sequences to fragments
- 6) comparison of the fragment sequences with a database
- 7) result: list of proteins from the database that have a similar fragmentation pattern





from: Gingras AC et al, 2004, J Physiol 563, 11-21

III) Protein Identification: Mass Spectrometry

- 3) mass spectrometry
- 4) comparison of the fragment sizes with a database


The peaks in the mass spectrum each represent a peptide present in the tryptic digest. A preliminary identification can be made by matching this spectrum to a database containing theoretical tryptic digests of all known proteins. This type of analysis is called peptide mass fingerprinting (PMF).



from: <http://www.med.monash.edu.au/biochem/facilities/proteomics/maldi.html>


III) Protein Identification: Mass Spectrometry

- 3a) further fragmentation of one of the fragments from the digest in the mass spectrometer, then again mass spectrometry (MSMS)
- 4a) comparison of the fragment sizes with a database





from: <http://www.med.monash.edu.au/biochem/facilities/proteomics/maldi.html>


III) Protein Identification: Mass Spectrometry



- 5) assignment of likely sequences to fragments
- 6) comparison of the fragment sequences with a database
- 7) result: list of proteins from the database that have a similar fragmentation pattern



III) Protein Identification: Mass Spectrometry



from: Gingras AC et al, 2004, J Physiol 563, 11-21


Limitations and artefacts

- contamination of gel bands with other proteins → identification of the contamination (very common: keratin from skin!) instead of the protein of interest
- not all proteins have the same detection efficiency (problems e.g. if many cysteines present) → even small contaminations sometimes lead to wrong identifications
- not all proteins are in the databases → database may show results that have a similar fragmentation pattern, but are otherwise unrelated

III) Protein Identification: N-terminal sequencing


Principle

Phenylisothiocyanate




Polypeptide

base



Phenylthiocarbamoyl-polypeptide

acid



Phenylthiohydantoin
derivative of the
N-terminal amino acid

identification of the
amino acid by
reversed phase HPLC
and comparison with
HPLC of standard
mixture

Polypeptide (N-1)

III) Protein Identification: N-terminal sequencing

Sample preparation

- 1) run acrylamide gel (denaturing or native)
- 2) blot onto PVDF (NOT nitrocellulose) membrane using glycine-free buffer
- 3) stain the membrane with ponceau red (CBB and other stains also work)
- 4) submit for sequencing

Limitations and artefacts


- problems with contaminations
- in eukaryotic proteins, the N-terminus is often blocked (e.g. by methylation), which required complicated de-blocking procedures. Also non-polymerised acrylamide remains in the gel can cause blocking, so let the gel polymerise over night
- Cys residues cannot be detected, and also glycosylated residues may appear as blank spaces in the sequence

IV) Protein Characterisation: Overview of Principles

- **Size determination**
- **Charge determination**
- **Analysis of cofactors**
- **Analysis of the 3-dimensional Structure**
 - **Activity tests**

IV) Protein Characterisation: Size determination

Comparison with predicted size by native and denaturing gel electrophoresis and by size exclusion chromatography can show native oligomerisation, post-translational modification but also artefactual degradation/aggregation




SDS gel and
Western blot of
TcHMA4

→ size shows
post-translational
processing as
cDNA sequence
predicts 128kDa

(Parameswaran,
Leitenmaier et al.,
2007, BBRC)


IV) Protein Characterisation: Charge determination

- Isoelectric focussing (IEF) can reveal the isoelectric point where the net charge of the protein is zero, i.e. balance between protonation of carboxyl groups and deprotonation of amino groups is achieved



IV) Protein Characterisation: Analysis of cofactors

- Identification and reactivity: UV/VIS Absorption and fluorescence Spectroscopy
- Metal content (about 30% of all proteins are metalloproteins!): AAS, ICP-MS/OES, EDX/PIXE/XRF




UV/VIS spectrum of TcHMA4

→ Cadmium binding causes ligand-metal-charge-transfer (LMCT) bands indicating cysteine ligation

Leitenmaier B, Witt A, Witzke A, Stemke A, Meyer-Klaucke W, Kroneck PMH, Küpper H (2011) Biochimica et Biophysica Acta - Biomembranes 1808, 2591-2599.

IV) Protein Characterisation: Analysis of cofactors

Metal ligands: EXAFS, EPR, heteronuclear NMR provide information about ligand types and their spatial arrangement around the active site (each of these techniques has different strenghts)




EXAFS spectrum of Zn-TcHMA4

→ mixed S/N ligands, multiple scattering shows histidine contribution

(Leitenmaier B, Witt A, Witzke A, Stemke A, Meyer-Klaucke W, Kroneck PMH, Küpper H (2011) Biochimica et Biophysica Acta - Biomembranes 1808, 2591-2599.)

IV) Protein Characterisation: Analysis of the 3D Structure

Circular Dichroism (CD) Spectroscopy




From: www.proteinchemist.com/cd/cdspec.html

- information about proportions of secondary structure types in a protein
- particularly useful when X-ray crystallography and NMR are not applicable

IV) Protein Characterisation: Analysis of the 3D Structure

X-ray crystallography

Example: in the nickel-binding transcription factor NikR, the mechanism was concluded from the X-ray structure




From: Schreiter ER et al., 2006, PNAS103, 13676-81

- Ni binding causes RHH domains to rotate about the flexible interdomain linkers to orient their antiparallel α -strands toward the same face of the repressor, allowing each to occupy the DNA major groove of an operator palindrome half-site
- Binding of Ni creates a surface of the MBD suitable for interacting with DNA by stabilization of a helix and a loop

IV) Protein Characterisation: Activity tests


Titration of an enzyme with its substrate(s) reveals binding constants and possible substrate inhibition



Activation of a human Cu-ATPase

Hung et al. Biochem.J. 2007, 401

Both are P_{1B} -Type ATPases showing the same activation pattern by „their“ metal after reconstitution into artificial lipid vesicles




Activity of TcHMA4

(Leitenmaier B, Witt A, Witzke A, Stemke A, Meyer-Klaucke W, Kroneck PMH, Küpper H (2011) Biochimica et Biophysica Acta - Biomembranes 1808, 2591-2599)

IV) Protein Characterisation: Activity tests

Two-dimensional data, e.g. substrate concentration and temperature reveal insights into interactions between factors that are decisive for enzyme activity, e.g. temperature dependence of substrate binding and influence of the substrate on the thermostability of the enzyme



(Leitenmaier B, Witt A, Witzke A, Stemke A, Meyer-Klaucke W, Kroneck PMH, Küpper H (2011) Biochimica et Biophysica Acta - Biomembranes 1808, 2591-2599)

Example: Flowchart of expression, isolation, purification and characterisation of the Cd/Zn-ATPase TcHMA4

Grow plants expressing TcHMA4 hydroponically, harvest and freeze roots

Grind the frozen roots with isolation buffer in liquid N₂, thaw

Centrifuge, discard the supernatant (soluble proteins) and resuspend the pellet (membrane proteins) with solubilisation buffer

Centrifuge for removing insoluble residue, collect solubilised protein

Immobilized Metal Affinity Chromatography on Ni-IDA column

Identify:
Western-Blotting,
Edman Sequ.

Quantify: In SDS
gel via fluorescent
dye

Metal binding:
AAS/ICP, EXAFS,
UV/VIS

Activity tests
(→ catalytical
properties)

**All slides of my lectures can be downloaded
from my workgroup homepage**

Biology Centre CAS → Institute of Plant Molecular Biology → Departments
→ Department of Plant Biophysics and Biochemistry,
or directly

http://webserver.umbr.cas.cz/~kupper/AG_Kuepper_Homepage.html